

# APOQUEL<sup>®</sup> DOSING GUIDE

- The dose of APOQUEL tablets is 0.4 up to 0.6 mg/kg
- APOQUEL should be given twice daily (BID) for up to 14 days of therapy and once daily (SID) thereafter for maintenance
- APOQUEL may be given with or without food

Weight Range (kg)	Number of Tablets to be Administered		
	3.6 mg	5.4 mg	16 mg
3.0 - 4.4	0.5		
4.5 - 5.9		0.5	
6.0 - 8.9	1.0		
9.0 - 13.4		1.0	
13.5 - 19.9			0.5
20.0 - 26.9		2.0	
27.0 - 39.9			1.0
40.0 - 54.9			1.5
55.0 - 80.0			2.0

[www.excellencein dermatology.co.nz](http://www.excellencein dermatology.co.nz)

zoetis

**apoquel**<sup>®</sup>  
(oclacitinib tablet)

Please refer to product labelling for full indications, precautions and contraindications.

Zoetis New Zealand Limited. Tel: 0800 963 847; [www.zoetis.co.nz](http://www.zoetis.co.nz). APOQUEL is a registered trade mark of Zoetis Inc. or its subsidiaries. ACVM No. A10963. RVM; Available only under Veterinary Authorisation. CA168