## LEPTOSPIROSIS VACCINATION

SPRING BORN CALVES												
	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	ОСТ	NOV	DEC
YEAR 1								BIRTH				
								VACC 1 – FROM 4 WEEKS OF AGE				
									VACC 2 4 – 6 WEEKS LATER*			
YEAR 2			AUTUMN BOOSTER*									
YEAR 3+			ANNUAL DRY OFF/ AUTUMN BOOSTER									

AUTUMN BORN CALVES												
	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
				BIRTH								
YEAR 1				VACC	1 – FROM	4 WEEKS OI	F AGE					
					VAC	CC 2 4-6	WEEKS LAT	ER#				
YEAR 2		AUTUMN BOOSTER										
YEAR 3+	YEAR 3+ ANNUAL DRY OFF/ AUTUMN BOOSTER											

<sup>\*</sup>Spring born calves require an autumn booster to align them with the herd and avoid long re-vaccination intervals.

<sup>\*</sup>Any calves receiving their second vaccine before 12 weeks of age will require a third vaccination around 6 months of age to manage the potential risk of MDA interference.







