# **Dosage and Administration:**

1mL/35 kg bodyweight by subcutaneous or intravenous injection 500kg cow dose = 14.5mL

#### Withholding periods:

Milk: Nil

Meat: 28 days

Rimadyl LA fast & effective relief from inflammation, fever & pain




For more information contact your veterinarian.



A division of Pfizer New Zealand Limited. Level 3, Pfizer House, 14 Normanby Road, Mt Eden, Auckland 1024, New Zealand. Tel: 0800 650 277, Fax: 0800 628 629. RIMADYL is a registered trademark of Pfizer Inc. or its subsidiaries. Registered pursuant to the ACVM Act 1997, No. A9897. Restricted Veterinary Medicine; Available only under Veterinary Authorisation. AM753002.

# Rimadyl® LA Effective relief from inflammation, fever and pain



# Rimadyl LA — Long acting Nil milk withhold.

# Inflammation, fever and pain occur in many common conditions including:

- Difficult calvings
- Downer cows
- Lameness
- Mastitis
- Metritis/RFMs (Retained Foetal Membranes)
- Pneumonia
- Surgical interventions, dehorning/ disbudding, castration, tailing
- Trauma and Injuries

The inflammation, fever and pain can decrease production by delaying healing, decreasing food and water intake and impairing immune function. Non steroidal anti-inflammatory drugs — NSAIDs for short

can help prevent this happening.

#### They work in 3 ways:

- Anti-inflammatory; meaning they reduce inflammation
- Anti-pyretic; meaning they reduce body temperature to a normal level
- Analgesic; meaning they reduce or alleviate pain


# Rimadyl LA is the NSAID we recommend because it:

#### Is fast and effective

- Reduces inflammation, which helps to prevent permanent tissue damage
- Reduces fever and heart rate, indicators of pain
- Aids in a rapid return to normal rumen function

# Is long acting

Just one injection and the cow gets maximum benefit from the prolonged treatment effect.



### Has nil milk withholding

This means there are no residue issues and treated cows can remain in supply, meaning less worry and more money in your pocket. Even though it has nil milk withholding, it is effective at treating the site of inflammation in mastitis - the udder tissue.