



# Ultravac™ Sheep & Beef Annual Treatment Programme

## ULTRAVAC™ 5in1 TREATMENT GUIDE

| SHEEP                       | 1mL DOSE                    | JAN   | FEB | MAR | APR | MAY | JUN                               | JUL | AUG   | SEP | OCT  | NOV | DEC |
|-----------------------------|-----------------------------|---|-----|-----|-----|-----|-----------------------------------|-----|---|-----|--|-----|-----|
| EWES                        | Previously Vaccinated       |   |     |     |     |     | Annual Booster 2–4 weeks pre-lamb |     |   |     |  |     |     |
|                             | Unknown Vaccination History | 1st dose at mating  |     |     |     |     | 2nd dose 2–4 weeks pre-lamb       |     |   |     |  |     |     |
| RAMS                        | Previously Vaccinated       | Annual Booster  |     |     |     |     |                                   |     |   |     |  |     |     |
|                             | Unknown Vaccination History | Two doses 4–6 weeks apart   |     |     |     |     |                                   |     |   |     |  |     |     |
| LAMBS                       | From Vaccinated Ewes        |   |     |     |     |     |                                   |     | 2 doses 4–6 weeks apart (from tailing or weaning)   |     |  |     |     |
|                             | From Unvaccinated Ewes      |   |     |     |     |     |                                   |     | PK/Antitet* at tailing<br>Followed by Ultravac 5in1 at weaning plus booster 4–6 weeks later |     |  |     |     |
| HOGGETS                     |                             | Ensure hoggets have received 2 doses (or annual booster) before autumn flush and winter crops |     |     |     |     |                                   |     |   |     |  |     |     |
| CATTLE                      | 2mL DOSE                    | JAN   | FEB | MAR | APR | MAY | JUN                               | JUL | AUG   | SEP | OCT  | NOV | DEC |
| ADULT COWS, HEIFERS & BULLS | Previously Vaccinated       | Annual Booster  |     |     |     |     |                                   |     |   |     |  |     |     |
|                             | Unknown Vaccination History | Two doses 4–6 weeks apart then normal annual booster  |     |     |     |     |                                   |     |   |     |  |     |     |
| CALVES                      |                             |   |     |     |     |     |                                   |     |   |     | 2 doses 4–6 weeks apart (as early as convenient) |     |     |

## ULTRAVAC™ 7in1 TREATMENT GUIDE

| CATTLE                      | 2.5mL DOSE                  | JAN  | FEB | MAR | APR | MAY | JUN | JUL | AUG | SEP | OCT   | NOV | DEC |
|-----------------------------|-----------------------------|--|-----|-----|-----|-----|-----|-----|-----|-----|---|-----|-----|
| ADULT COWS, HEIFERS & BULLS | Previously Vaccinated       | Annual Booster (recommended in Autumn)               |     |     |     |     |     |     |     |     |   |     |     |
|                             | Unknown Vaccination History | Two doses 4–6 weeks apart then normal annual booster |     |     |     |     |     |     |     |     |   |     |     |
| CALVES                      |                             |  |     |     |     |     |     |     |     |     | 2 doses 4–6 weeks apart (from 4 weeks of age) |     |     |

PK/Antitet also referred to as Lamb Vaccine-provides immediate short term protection against tetanus & is a priming vaccination for pulpy kidney only.



A division of Pfizer New Zealand Ltd, Level 3, Pfizer House, 14 Normanby Road, Mt Eden, Auckland. Ultravac™ 5in1 and Ultravac™ 5in1 with Selenium are trademarks of Pfizer. Registered pursuant to the ACVM Act 1997, Nos. A3585, A6926. OTC. Ultravac™ 7in1 is a trademark of Pfizer. Registered pursuant to the ACVM Act 1997 No. A6935. Prescription Animal Remedy (PAR) Class I. For use only under the authority or prescription of a veterinarian. AM029033. H&T PAH0565.

