

Effective mastitis solutions for every situation

	CLAVULOX L.C.		ORBENIN L.A.		MASTALONE
Time of Season	Calving		Lactation/ Dry Period		All season
Treatment Regime	3 tubes at 12 hour intervals	5 tubes at 12 hour intervals	3 or 5* tubes at 48 hour intervals	5 tubes at 24 hour intervals	3 tubes at 24 hour intervals
Milk withholding period twice a day	48 hours/ 4 milkings	96 hours/ 8 milkings	84 hours/ 7 milkings	96 hours/ 8 milkings	96 hours/ 8 milkings
Milk withholding period once a day			72 hours/ 3 milkings	96 hours/ 4 milkings	
Meat withholding period	3 days	7 days	3 days		30 days
Effective against penicillin resistant Staphs	✓		✓		✓
Anti-inflammatory	✓				✓

*Recommended for treatment of subclinical mastitis during lactation which require longer treatment regimes.

Zoetis Mastitis Solutions

LACTATING COW TREATMENTS



CLAVULOX L.C.



ORBENIN L.A.



MASTALONE

DRY COW PRODUCTS



TEATSEAL



ORBENIN ENDURO



ORBENIN DC

www.dairywellness.co.nz

Use only as directed by your veterinarian.
Use antibiotics responsibly.

Zoetis New Zealand Limited. Tel: 0800 963 847; www.zoetis.co.nz. ORBENIN and CLAVULOX are registered trade marks of GlaxoSmithKline NZ Ltd. ACVM No. A3664, A5945, A0888 and A6036. MASTALONE and TEATSEAL are registered trade marks of Zoetis Inc. ACVM A0829 and A7294. RVM; Available only under Veterinary Authorisation.

zoetis

Effective mastitis solutions for every situation



zoetis

Several factors must be considered when choosing a mastitis treatment including:

- **Time of season**
- **Aetiology**
- **Severity of mastitis**
- **Initial case or repeat**
- **Milking frequency**
- **Withholding periods**

Milk quality consultations are an ideal time to review herd mastitis patterns and treatments with dairy farmers.

Mastitis treatments should be chosen based on expected aetiology; cow factors such as severity and stage of lactation; and farm factors such as prevalence and milking frequency. The three Zoetis lactating cow products each have unique properties, allowing veterinarians to prescribe mastitis treatments more specifically to achieve the highest chances of cure the first time, while minimising time out of the vat.

Many veterinarians prescribe according to stage of lactation or time of year. The Zoetis range covers the whole lactation. The table matches the product's key benefits with the most appropriate time to use it.

	Calving A broad spectrum antibiotic to treat predominantly <i>Strep. uberis</i> with some <i>Staph. aureus</i> and coliforms.	Lactation An antibiotic to treat predominantly <i>Staph. aureus</i> with some <i>Strep spp.</i>	Tough Cases An antibiotic to treat difficult cases of mastitis where other treatments have failed. Predominantly <i>Staph. aureus</i> with some <i>Strep spp.</i>
CLAVULOX® L.C.			
ORBENIN® L.A.			
MASTALONE®			

CLAVULOX® L.C.

The Spring Specialist

Broad spectrum with high efficacy against *Strep. uberis* making it ideal for first line treatments around calving

Contains the anti-inflammatory prednisolone

Short milk and meat withholding periods

Contains the antibiotic amoxicillin potentiated with clavulanic acid



ORBENIN® L.A.

The Workhorse

Highly effective at treating *Staph. aureus*

A unique long acting base allows a 48 hour treatment interval

Registered for Once A Day milking

Short meat withholding period

Contains the antibiotic cloxacillin



MASTALONE®

The Big Gun

An effective treatment for severe mastitis or as a second line treatment where others have failed

Contains the anti-inflammatory prednisolone

24 hour treatment interval

Contains the antibiotics oxytetracycline, oleandomycin and neomycin



ZOETIS TEATWIPES

Use for Best Practice Administration

99% BACTERICIDAL EFFECT*.

Made from high quality textured cloth.

Easy dispensing container to save time.



* Independently tested by New Zealand Laboratory Services.